

Lattice analysis of semi-leptonic form factors*

Tanmoy Bhattacharya and Rajan Gupta ^a

^aT-8 Group, MS B285, Los Alamos National Laboratory, Los Alamos, New Mexico 87545 U. S. A.

We present preliminary results from simulations done on 170 $32^3 \times 64$ lattices at $\beta = 6.0$ using quenched Wilson fermions. This talk focuses on the Q^2 behavior of the form-factors, extrapolation in quark masses, dependence on renormalization scheme, and comparison with heavy-quark effective theory (HQET). Even though we cannot estimate errors due to quenching and discretization, our results are consistent with experimental results for D decays. We present results for the Isgur-Wise function and estimate $\xi'(w=1) = 0.97(6)$.

1. TECHNICAL DETAILS


We briefly mention some details of our analysis to extract the form-factors at $Q^2 = 0$ relevant to phenomenology. A full analysis will be presented elsewhere [1]. The details of the lattices are contained in the papers on the hadron spectrum [2] [3], and decay constants [4]. Preliminary results on a sub-set of lattices have been presented at LATTICE 94 [5] and DPF 94 [6]. We do not have data to extrapolate to m_b or to $a = 0$, thus our results are relevant for D decays, *i.e.* $D \rightarrow K l \nu$, $D \rightarrow \pi l \nu$, $D \rightarrow K^* l \nu$, and $D \rightarrow \rho l \nu$, calculated at $a^{-1} = 2.33(4)$ GeV.

The decaying D meson is created at rest by using a $\vec{p} = 0$ source. On each lattice we make two measurements by creating the D meson at $t = 7$ and 57, for a total statistical sample size of 340. The sink for the final state meson is at $t = 32$ in both cases. The time-slice of the weak operator is taken to vary between $10 \leq t \leq 30$ and $35 \leq t \leq 55$ respectively, and the insertion is at 5 lowest values of lattice momenta. The quark propagators are created using a Wuppertal smeared source as described in [2].

In order to isolate the desired matrix element (ME) we construct a ratio of 3-point to 2-point correlation functions. We have a choice of using either smeared-smeared (SS) or smeared-local (SL) 2-point correlation functions. We calculate

*Based on talks presented by Tanmoy Bhattacharya and Rajan Gupta. These calculations have been done on the CM5 at LANL as part of the DOE HPCC Grand Challenge program, and at NCSA under a Metacenter allocation.

Figure 1. Ratio of 3-point to 2-point SL function versus the time slice of the operator insertion.




the ME both ways and take the average as our best estimate. Figure 1 shows a typical example of signal for the ratio of correlators in the SL case: the quality in the SS case is very similar. We have seen a steady improvement in the consistency between these two estimates of ME with statistics. With the current sample they are within 1σ in all cases.

Pole dominance hypothesis (PDH): It states that all form-factors, $f(Q^2)$, have the structure

$$f(Q^2) = f(0)/(1 - Q^2/M^2), \quad (1)$$

where M is the mass of the nearest resonance

Figure 2. Three fits to test Q^2 behavior of f_+ .


with the right quantum numbers. To test PDH we make two kinds of fits: (i) single parameter “pole” fit where M is the lattice measured value of the resonance mass, (ii) two parameter “best” fit where M and $f(0)$ are free parameters. An example of these fits is shown in figure 2. Overall we find that only f_0 and f_V are well described by the “pole” form. f_+ and f_{A_0} are consistent with “pole” form with $M < M_{pole}$, while for f_{A_3} we find $M > M_{pole}$. f_{A_1} and f_{A_2} show a much smaller Q^2 dependence than expected from pole dominance, however the data are too noisy to make a definite statement. We use results from “best” fit for our final estimates.

HQET: At leading order in $1/m_c$, HQET predicts that the Isgur-Wise function ξ describes all form-factors relevant to $D \rightarrow K l \nu$ and $D \rightarrow K^* l \nu$ decays. Neglecting $O(\alpha_s)$ corrections, one gets [7]

$$\begin{aligned} \xi(w) &= R f_+(q^2) = R \left(1 - \frac{q^2}{(M_i + M_f)^2}\right)^{-1} f_0(q^2) \\ &= R^* V(q^2) = R^* A_0(q^2) = R^* A_2(q^2) \\ &= R^* \left(1 - \frac{q^2}{(M_i + M_f)^2}\right)^{-1} A_1(q^2), \end{aligned}$$

where M_i and M_f are the initial and final meson masses and

$$w = v_i \cdot v_f = \frac{M_i^2 + M_f^2 - q^2}{2M_i M_f}; \quad R = \frac{2\sqrt{M_i M_f}}{M_i + M_f}.$$

Figure 3. Comparison of $f_+(q^2)$ (squares) and $(1 - \frac{q^2}{(M_i + M_f)^2})^{-1} f_0(q^2)$ (diamonds) versus w .

The first HQET relation, $(1 - q^2/(M_i + M_f)^2)^{-1} f_0 = f_+$, is satisfied by our data as exemplified in figure 3 for $CU_3 \rightarrow U_1 U_3$ decay. Thus, both f_0 and f_+ cannot simultaneously satisfy the PDH. If leading order HQET holds, *i.e.* $M_i + M_f \sim m_c + m_s \sim M_{pole}$, f_+ should obey a ‘dipole’ form as f_0 agrees with pole dominance. The data, as shown in figure 2, suggests that f_+ lies in between “pole” and “dipole” forms. Similar analysis for the vector form factors is under progress.

Dependence on quark mass: Figure 4 shows an example of the variation of $f_+(0)$ with quark masses. There is significant dependence on the mass of the quark C decays into. This is a kinematic effect as shown in Section 4. Our data is good enough to expose slight dependence on $m_{spectator}$ —the small decrease in slope between the transitions $CU_i \rightarrow U_1 U_i$ and $CU_i \rightarrow SU_i$ is consistent with HQET.

Fixing m_s : We fix m_s using M_K^2/M_ρ^2 and M_ϕ/M_ρ . Our preferred way is $m_s(M_\phi)$, and the variation of results with m_s is shown in Table 2.

$O(ma)$ effects: In Ref. [2] we discuss $O(ma)$ effects for heavy-light mesons. The kinetic mass $M_2 \equiv d^2 E/dp^2$ is given by $\sinh M$ as the data agree with the dispersion relation $\sinh^2 E/2 =$

Figure 4. Extrapolation of $f_+(Q^2 = 0)$ to m_u . $f_+(D \rightarrow Kl\nu)$ is extracted from points labeled by squares and octagons, while $f_+(D \rightarrow \pi l\nu)$ is from data with degenerate $q\bar{q}$ points (crosses).




Table 1. Estimates of form factors in 3 commonly used renormalization schemes defined in [4].

	TAD1	TAD π	TADU $_0$
f_+	0.71(4)	0.75(4)	0.66(4)
f_0	0.73(3)	0.77(3)	0.68(2)
f_V	1.28(7)	1.36(7)	1.19(6)
f_{A_0}	0.84(3)	0.85(3)	0.79(3)
f_{A_1}	0.72(3)	0.74(3)	0.68(3)
f_{A_2}	0.49(9)	0.50(9)	0.46(8)
f_{A_3}	0.85(3)	0.87(3)	0.80(3)
f_V/f_{A_1}	1.78(7)	1.84(8)	1.76(7)

$\sin^2 p/2 + \sinh^2 M/2$. We show variation of form factors with M in Table 2.

Renormalization Constants: To relate lattice results to experimental data we need the renormalization constants Z_A and Z_V . We use three Lepage-Mackenzie tadpole improved schemes described in Ref. [4]. Our preferred scheme is *TAD1*, and the variation with the schemes is illustrated in Table 1.

2. RESULTS at $\beta = 6.0$

Our final results for $D \rightarrow Kl\nu$ and $D \rightarrow K^*l\nu$ are shown in tables 1 and 2 along with variation with m_s , type of fit, heavy-light meson

Table 2. Estimates in *TAD1* scheme at $Q^2 = 0$. The variations give estimates of systematic errors.

		Pole		Best	
		$m_s(M_K)$	$m_s(M_\phi)$	$m_s(M_K)$	$m_s(M_\phi)$
f_+	M_1	0.78(2)	0.80(2)	0.70(4)	0.71(4)
	M_2	0.79(2)	0.80(2)	0.67(5)	0.68(4)
f_0	M_1	0.70(2)	0.72(2)	0.71(3)	0.73(3)
	M_2	0.72(2)	0.73(2)	0.68(3)	0.70(2)
f_V	M_1	1.27(7)	1.28(6)	1.27(7)	1.28(7)
	M_2	1.22(6)	1.23(6)	1.22(9)	1.23(9)
f_{A_0}	M_1	0.85(3)	0.85(3)	0.83(3)	0.84(3)
	M_2	0.84(3)	0.84(3)	0.80(3)	0.81(3)
f_{A_1}	M_1	0.67(2)	0.68(2)	0.71(3)	0.72(3)
	M_2	0.65(2)	0.66(2)	0.67(4)	0.69(4)
f_{A_2}	M_1	0.46(9)	0.47(8)	0.49(9)	0.49(9)
	M_2	0.42(9)	0.44(9)	0.50(14)	0.52(12)
f_{A_3}	M_1	0.84(3)	0.85(3)	0.84(3)	0.85(3)
	M_2	0.81(3)	0.82(3)	0.81(4)	0.82(4)
f_0/f_+	M_1	0.93(1)	0.94(1)	1.02(4)	1.02(3)
	M_2	0.95(1)	0.95(1)	1.03(4)	1.02(4)
f_V/f_{A_1}	M_1	1.83(7)	1.81(6)	1.79(8)	1.78(7)
	M_2	1.85(7)	1.83(6)	1.81(11)	1.79(10)
f_{A_2}/f_{A_1}	M_1	0.70(10)	0.70(9)	0.69(12)	0.68(11)
	M_2	0.65(11)	0.67(11)	0.74(18)	0.75(15)

mass, and the renormalization scheme. Our preferred estimates are with $m_s(M_\phi)$, “best” fit, M_1 , and *TAD1* scheme. We also get that $f_+^\pi/f_+^K = 0.87(4)$.


3. $d\Gamma(Q^2)$

As explained in section 1 there are considerable uncertainties involved in extrapolating the form factors to $Q^2 = 0$. Therefore, we also calculate $d\Gamma(Q^2)/dQ^2$ by linearly extrapolating in m_q the form-factors at fixed 3-momentum transfer. In figure 5 we show the results for $D \rightarrow Ke\bar{\nu}_e$. In figure 6 we show the longitudinal, transverse and the total decay widths for the process $D \rightarrow K^*e\bar{\nu}_e$.

4. Isgur-Wise function

As mentioned in section 1, at the leading order in the heavy quark mass, there is one universal ‘Isgur-Wise’ function ξ which controls all the form

Figure 5. $(1/V_{cs}^2)d\Gamma(q^2)/dq^2$ versus q^2 in GeV^2 . The shape is in qualitative agreement with experimental data.



factors. In figure 7 we present data for $\xi_{ren}(w)$,

$$h_+(w) = \left[\hat{C}_1(w) + \frac{1+w}{2}(\hat{C}_2(w) + \hat{C}_3(w)) \right] \xi_{ren}(w) \\ = f_+(q^2) \frac{M_i + M_f}{2\sqrt{M_i M_f}} + f_-(q^2) \frac{M_i - M_f}{2\sqrt{M_i M_f}},$$

where $\hat{C}_1, \hat{C}_2, \hat{C}_3$ are HQET renormalization constants including the leading $O(\alpha_s)$ corrections [7]. The data show a slight dependence on the spectator quark mass indicative of m_c^{-1} corrections. Analysis from the vector form factors is in progress. From this data we estimate the slope to be $\xi'(w=1) = 0.97(6)$.

REFERENCES

1. T. Bhattacharya, R. Gupta, in preparation.
2. T. Bhattacharya, et al., LA-UR-95-2354.
3. T. Bhattacharya, R. Gupta, and S. Sharpe, LA-UR-95-4224.
4. T. Bhattacharya, R. Gupta, LA-UR-95-2355.
5. T. Bhattacharya, R. Gupta, Nucl. Phys. B (Proc. Suppl.) 42 (1995) 935.
6. T. Bhattacharya, R. Gupta, DPF94 proceedings, World Scientific 1995, page 605.
7. M. Neubert, Phys. Rep. 245 (1994) 259.

Figure 6. $(1/V_{cs}^2)d\Gamma(q^2)/dq^2$ versus q^2 in GeV^2 .




Figure 7. $\xi(w)/\xi(1)$ at various quark masses and momenta. The symbols label the flavor that the C quark decays to, and variation of ξ with w is a kinematic effect. Dependence on $m_{spectator}$ is shown most clearly by the clusters of 3 points at $w \approx 1.2$. The largest value in each cluster corresponds to the lightest spectator (U_3). The data are fit to $\xi(w) = \frac{2}{w+1} \exp(-(2\rho^2 - 1)\frac{w-1}{w+1})$, from which we estimate the slope $\xi'(1)$.

